

Instructions: This was a 2 hour, closed notes, closed book, and pledged exam.

1. Find and classify all the critical points of

$$f(x, y) = \frac{1}{2}x^2 - xy + \frac{1}{3}y^3.$$

Solution: Since f is differentiable everywhere, the only critical points are the points where the gradient vanishes. $\nabla f(x, y) = (x - y, y^2 - x) = (0, 0)$ precisely at the points $(0, 0)$ and $(1, 1)$. Note $\frac{\partial^2 f}{\partial x^2} = 1$, $\frac{\partial^2 f}{\partial y \partial x} = -1$, and $\frac{\partial^2 f}{\partial y^2} = 2y$. Apply the second derivative test:

point	$\frac{\partial^2 f}{\partial x^2}$	D	classification
$(0, 0)$	1	-1	saddle
$(1, 1)$	1	1	strict local min

2. Let $g(x, y) = 2e^{-x} \cos y$.

- (a) Find the quadratic Taylor polynomial for $g(x, y)$ around the point $(0, 0)$.

$$\begin{aligned} g(0, 0) &= 2 \\ \frac{\partial g}{\partial x} \Big|_{(0,0)} &= -2e^{-x} \cos y \Big|_{(0,0)} = -2 \\ \frac{\partial^2 g}{\partial x^2} \Big|_{(0,0)} &= 2e^{-x} \cos y \Big|_{(0,0)} = 2 \\ \frac{\partial^2 g}{\partial y \partial x} \Big|_{(0,0)} &= -2e^{-x} \sin y \Big|_{(0,0)} = 0 \\ \frac{\partial g}{\partial y} \Big|_{(0,0)} &= -2e^{-x} \sin y \Big|_{(0,0)} = 0 \\ \frac{\partial^2 g}{\partial y^2} \Big|_{(0,0)} &= -2e^{-x} \cos y \Big|_{(0,0)} = -2 \end{aligned}$$

So the quadratic Taylor polynomial for $g(x, y)$ around $(0, 0)$ is

$$g(h_1, h_2) \approx 2 - 2h_1 + h_1^2 - h_2^2$$

- (b) Use your answer in part (a) to estimate $2e^{-0.2} \cos 0.4$.

We just evaluate $g(0.2, 0.4) \approx 2 - 2(0.2) + (0.2)^2 - (0.4)^2 = 2 - .4 + .04 - .16 = 1.48$.
(The actual value is approximately 1.508202).

3. A tank is in the shape of a half-cylinder of radius 2 and height 3. It is situated in \mathbb{R}^3 , given by the inequalities $\sqrt{x^2 + y^2} \leq 2$, $y \geq 0$, and $0 \leq z \leq 3$. The temperature at the point (x, y, z) is given by

$$T(x, y, z) = 2yz^2 \sqrt{x^2 + y^2} \text{ } ^\circ\text{C}.$$

Find the average temperature in the tank.

Solution: We describe the tank in cylindrical coordinates as $0 \leq r \leq 2, 0 \leq \theta \leq \pi$, and $0 \leq z \leq 3$. Recall the formula $[T]_{av} = \frac{\iiint_W T(x,y,z) dV}{\iiint_W dV}$.

We use cylindrical coordinates to compute

$$\begin{aligned} \iiint_W T(x,y,z) dV &= \int_0^3 \int_0^\pi \int_0^2 2(r \sin \theta)(z^2)(r) r dr d\theta dz \\ &= 2 \left(\int_0^3 z^2 dz \right) \left(\int_0^\pi \sin \theta d\theta \right) \left(\int_0^2 r^3 dr \right) \\ &= 2(2)(4)(9) = 144 \end{aligned}$$

The denominator (volume of region) is given by the formula $\text{Vol}(W) = \frac{\pi \cdot 2^2 \cdot 3}{2} = 6\pi$, or by computing

$$\iiint_W dV = \int_0^3 \int_0^\pi \int_0^2 r dr d\theta dz = \left(\int_0^3 dz \right) \left(\int_0^\pi d\theta \right) \left(\int_0^2 r dr \right) = (3)(\pi)(2) = 6\pi$$

Thus, $[T]_{av} = \frac{144}{6\pi} = \frac{24}{\pi} \circ C$.

4. Let T be the triangle with vertices $(0, 0)$, $(1, 1)$ and $(0, 1)$ and let $f(x, y) = x \sin(y^3)$.

(a) Find the correct limits of integration to **set up** $\iint_T f(x, y) dA$ as a double integral

$$\iint f(x, y) dx dy.$$

Solution: $\int_0^1 \int_0^y f(x, y) dx dy$.

(b) Find the correct limits of integration to **set up** $\iint_T f(x, y) dA$ as a double integral

$$\iint f(x, y) dy dx.$$

Solution: $\int_0^1 \int_x^1 f(x, y) dy dx$.

(c) Compute $\iint_T f(x, y) dA$.

Solution: Use the set up from (a):

$$\begin{aligned} \iint_T f(x, y) dA &= \int_0^1 \int_0^y x \sin(y^3) dx dy \\ &= \int_0^1 \frac{\sin(y^3)}{2} x^2 \Big|_{x=0}^y dy \\ &= \int_0^1 \frac{1}{2} y^2 \sin(y^3) dy \\ &= -\frac{\cos(y^3)}{6} \Big|_{y=0}^1 \\ &= \frac{1 - \cos 1}{6} \end{aligned}$$

5. Find the maximum and minimum values obtained by $f(x, y) = x + y^2$ on the ellipse $x^2 + 3y^2 \leq 9$.

Solution: First, find critical points in the interior $x^2 + 3y^2 < 9$. Note $\nabla f(x, y) = (1, 2y)$ is never $(0, 0)$, so there are no critical points in the interior.

Second, find critical points on the boundary $x^2 + 3y^2 = 9$ using Lagrange Multipliers. Our constraint function is $g(x, y) = x^2 + 3y^2$. Solve $\nabla f(x, y) = \lambda \nabla g(x, y)$, i.e. $(1, 2y) = \lambda(2x, 6y)$. The second coordinate gives two possibilities: $y = 0$ or $\lambda = 1/3$. If $y = 0$, then $x = \pm 3$ (from the constraint $x^2 + 3y^2 = 9$). If $\lambda = 1/3$, then $x = 3/2$ (from $1 = 2\lambda x$), and the constraint gives $y = \pm 3/2$. There are four critical points to investigate: $(\pm 3, 0)$ and $(3/2, \pm 3/2)$.

(x, y)	$f(x, y)$
$(3, 0)$	3
$(-3, 0)$	-3
$(3/2, 3/2)$	15/4
$(3/2, -3/2)$	15/4

Thus the absolute maximum value of f on boundary is $15/4$, and the absolute minimum value on the boundary is -3 .

Since there are no critical points from the interior, these maximum and minimum boundary values are also the maximum and minimum values throughout the entire region.

6. The region S is cut from a solid ball of radius 1 centered at the origin. S is the region cut by the inequalities $z \geq 0$ and $y \geq x$. (S is one-quarter of the entire ball, and contains the point $(0, 1, 0)$.)

The mass density of S at a point (x, y, z) is given by the function $\delta(x, y, z) = 30z^2 \text{ kg/m}^3$.

- (a) Find the total mass of S .

Solution: The total mass is given by $\iiint_S \delta(x, y, z) dV$. Note that S is described in spherical coordinates by $0 \leq \rho \leq 1$, $\pi/4 \leq \theta \leq 5\pi/4$, and $0 \leq \phi \leq \pi/2$. Thus

$$\begin{aligned}
 \iiint_S \delta(x, y, z) dV &= \int_0^{\pi/2} \int_{\pi/4}^{5\pi/4} \int_0^1 30(\rho \cos \phi)^2 \rho^2 \sin \phi d\rho d\theta d\phi \\
 &= 30 \left(\int_0^{\pi/2} \cos^2 \phi \sin \phi d\phi \right) \left(\int_{\pi/4}^{5\pi/4} d\theta \right) \left(\int_0^1 \rho^4 d\rho \right) \\
 &= 30 \left(\left. \frac{-\cos^3 \phi}{3} \right|_0^{\pi/2} \right) (\pi) \left(\frac{1}{5} \right) \\
 &= 2\pi \text{ kg}
 \end{aligned}$$

- (b) Find the average mass density of S .

Solution: Average mass density is

$$[\delta]_{\text{av}} = \frac{\iiint_S \delta(x, y, z) dV}{\iiint_S dV} = \frac{2\pi \text{ kg}}{\text{Vol}(S)} = \frac{2\pi \text{ kg}}{\pi/3 \text{ m}^3} = 6 \text{ kg/m}^3$$