

Subgroups of the mapping class  
group and higher-order  
signature cocycles

Shelly Harvey (Rice University)  
joint with Tim Cochran (Rice University)  
+ Peter Horn (Rice University)

Goal: For certain subgroups  $J(H)$  of the mapping class group of a surface, define

- quasi-homomorphisms

$$\rho^\Psi : J(H) \longrightarrow \mathbb{R}$$

- 2-cocycles


$$\sigma_\Psi : J(H) \times J(H) \longrightarrow \mathbb{R}.$$

When  $J(H) = \text{Torelli subgroup}$

and  $\Psi: H_1(\Sigma) \rightarrow U(1)$ , we will show

that  $\{\rho^4\}$  has infinite rank.

Let  $\Sigma = \Sigma_{g,1}$  be a compact, oriented surface with 1 boundary component.




$F :=$  free group generated by  $\{x_1, \dots, x_{2g}\}$   
 $\cong \pi_1(\Sigma, x_0),$   
 $:= \{\text{loops in } \Sigma \text{ based at } x_0\}/\text{homotopy}$


Def: The mapping class group of  $\Sigma$ ,  $\text{Mod}(\Sigma)$ , is the group of isotopy classes of orientation preserving homeomorphisms of  $\Sigma$  that fix the boundary of  $\Sigma$  pointwise.

Ex:

(for a closed surface)



Hyperelliptic involution




$\text{Mod}(\Sigma)$  is related to 3-manifolds

(Assume  $\Sigma \neq \emptyset$ ). Let  $f: \Sigma \rightarrow \Sigma$ .

1. let  $U_i$  be a thickened graph  $\infty$


Glue  $U_1$  to  $U_2$

using  $f: \partial U_1 \rightarrow \partial U_2$



Get a Heegaard decomposition of  $M = U_1 \cup_f U_2$

$$2. M_f = \Sigma \times I / (x, 1) \sim (f(x), 0)$$




$f$

glue top to  
bottom by  
 $f$


$M_f$  is a fiber bundle over  $S^1$  with fiber  $\Sigma$ .

## Dehn twists



Let  $\alpha$  be a S.C.C. on  $\Sigma$ . Let  $A = \text{nbhd}$  of  $\gamma$ .




Define  $D_\alpha : \Sigma \rightarrow \Sigma$  by  $\bullet D_\alpha|_{\Sigma - A} = \text{id}$




Ex:  $f = D_B \circ D_\alpha$





$D_\alpha$




$D_B$




Ex:



$\downarrow D_\alpha$



Theorem (Dehn, 1922)  $\text{Mod}(\Sigma)$  is generated by finitely many Dehn twists.



Theorem (McCool, Hatcher-Thurston, Deligne-Mumford):

For any surface  $\text{Mod}(\Sigma)$  is finitely presented.

- Hatcher-Thurston found an algorithm to construct a presentation. Harer produced the first explicit presentation.

Let  $f \in \text{Mod}(\Sigma)$   $\Rightarrow f|_{\partial\Sigma} = \text{id}$  so by

choosing a base point  $x_0 \in \partial\Sigma$ ,

$f_* : \pi_1(\Sigma, x_0) \rightarrow \pi_1(\Sigma, x_0)$  is an isomorphism.

Theorem (Dehn-Nielsen-Baer) :

$$\tau : \text{Mod}(\Sigma) \hookrightarrow \text{Aut}(\pi_1(\Sigma, x_0)) = \text{Aut}(F)$$

$$(f : \Sigma \rightarrow \Sigma) \longmapsto (f_* : \pi_1(\Sigma) \rightarrow \pi_1(\Sigma))$$

is a monomorphism. Hence  $f = g$

$$\Leftrightarrow f_* = g_* !$$

$$\mathcal{I} : \text{Mod}(\Sigma) \hookrightarrow \text{Aut}(\pi_1(\Sigma, x_0)) = \text{Aut}(F)$$

$$f \longmapsto f_*$$

Can approximate  $\mathcal{I}$  by send  $f \mapsto f_*^k$   
 where  $f_*^k : F/F_k \longrightarrow F/F_k$  for  $\{F_k\}$

a characteristic series that "approximates  $F$ ."

$$\mathcal{I}_k : \text{Mod}(\Sigma) \longrightarrow \text{Aut}(F/F_k)$$

$$f \longmapsto f_*^k$$

Example:  $F_2 = [F, F]$   
 $= \text{Subgp gen by } aba^{-1}b^{-1} \text{ where}$   
 $a, b \in F$


Def: The Torelli group of  $\Sigma$ ,  $\mathcal{I}(\Sigma)$  is the subgroup of  $\text{Mod}(\Sigma)$  consisting of diffeos that induces the identity on  $H_1(\Sigma) = F/F_2$ .

$$\mathcal{I}(\Sigma) = \ker(\tau_2 : \text{Mod}(\Sigma) \rightarrow \text{Aut}(H_1(\Sigma)))$$

Thm (Powell, Birman, Johnson):  $I(\Sigma)$  generated by

Dehn twists on bounding pairs of curves,

$$D_\alpha \circ D_\beta^{-1}.$$



$\alpha, \beta$  disjoint  
homologous  
curves

Thm (Johnson):  $I(\Sigma)$  is finitely generated for  $g \geq 3$ .

Thm (Mess):  $I(\Sigma)$  is an infinitely generated free group when  $g = 2$ .

We wish to investigate the homology and cohomology of certain special subgroups of  $I(\Sigma)$  using  $\rho$ -invariants of 3 and 4-dimensional manifolds.

Def: Let  $G$  be a group. Then the lower-central series of  $G$  is defined

as


- $G_2 := [G, G] =$  commutator subgp of  $G$ .
- for  $k \geq 3$ ,  $G_{k+1} := [G_k, G]$ .

Def: The  $k^{\text{th}}$  generalized Johnson subgroup of  $\Sigma$ ,  $J_k = J_k(\Sigma)$  is the subgroup of  $\text{Mod}(\Sigma)$  consisting of homeos that induce the identity on  $F/F_k$ .

- Note that  $J_2(\Sigma) = I(\Sigma)$ .

Thm (Johnson):  $J_3(\Sigma)$  = subgroup generated by Dehn twists on separating simple closed curves on  $\Sigma$ .

$\alpha$  is separating



$[J_3(\Sigma)$ , denoted  $K_g$ , is often called the  
Johnson subgroup.]

Question (Morita): Is  $H_1(K_g; \mathbb{Z})$ , the abelianization of  $K_g$ , finitely generated for  $g \geq 2$ ? [When  $g=2$ ,  $I_2 = K_2 = \text{infinitely gen. free gp}$ ].

Note: Still unknown if  $K_g$  (or more generally  $J_k(\Sigma), k \geq 3$ ) is finitely generated for  $g \geq 3$ !

## Construction of Invariants

Let  $H \trianglelefteq F = \pi_1(\Sigma)$ , a characteristic subgroup, and  $\underline{J(H)} =$  subgroup of  $\text{Mod}(\Sigma)$  consisting of homeos of  $\Sigma$  that induce the identity on  $F/H$ .

e.g.  $H \subseteq F_K \Rightarrow J(H) \subseteq K^{\text{th}}$  Johnson subgroup.

Given  $f \in \mathcal{J}(H) \rightsquigarrow$  closed 3-mfld  $N_f$  by


- $M_f = \text{mapping torus}$   
of  $f$

$$= \Sigma \times I / (x, 1) \sim (f(x), 1)$$


$\wedge x \in \Sigma$



- $N_f = M_f / (y, t) \sim (y, s)$   
for  $t, s \in I$   
and  $y \in \partial \Sigma$



$\alpha$  identified to  
a point in  $N_f$ .



Then  $\pi_1(M_f) = \langle x_1, \dots, x_{2g}, t \mid tx_i t^{-1} = f(x_i) \rangle$

$\downarrow$  kill  $[x] = t$

$$\pi_1(N_f) = \langle x_1, \dots, x_{2g} \mid x_i = f(x_i) \rangle$$

but since  $f(x_i) \equiv x_i \pmod{H} \Rightarrow$

$$\pi_1(N_f)/H \cong F/H$$

Thus, we have  $\Psi_H: \pi_1(N_f) \rightarrow F/H$ .

↑  
independent of  $f$

## Cheeger-Gromov $\rho$ -invt

If  $M^3$  is a closed, oriented 3-mfld,  
and  $\Psi: \pi_1 M \rightarrow \Gamma = \text{countable gp}$ ,  
Cheeger-Gromov (generalizing Atiyah-  
Patodi-Singer) define the topological  
invariant :

$$\rho^{(2)}(M, \Psi: \pi_1 M \rightarrow \Gamma) \in \mathbb{R}$$

Def:  $\rho_H(f) = \rho^{(2)}(N_f, \psi_H: \pi_1(N_H) \rightarrow F/H) \in \mathbb{R}$ ,

the Cheeger Gromov  $L^2$ - $\rho$  invariant  
associated to  $(N_f, \psi_H)$ .

Note:  $L^2$ - $\rho$  invariants have been useful recently  
in the study of 3-manifolds and  
knot and link concordance (Cochran, H, Leidy,  
Kim, Cha, Orr, Teichner, P. Horn, P. Heck)

Lemma For each  $(M^3, \psi)$  there exists  $(W^4, \phi)$

s.t.

$$\begin{array}{ccc} \pi_{1,2} W = \pi_1 M & \xrightarrow{\psi} & \Gamma \\ \downarrow & & \downarrow \\ \pi_1 W & \xrightarrow{\phi} & \Lambda \end{array}$$

Thm (Ramachandran)

$$Q^{(2)}(M, \Gamma) = \left( \begin{array}{l} L^2\text{-signature of } \Lambda\text{-equivariant} \\ \text{intersection form on } H_2(W, \mathbb{Z}\Lambda) \end{array} \right)$$

- (signature of  $W$ ).  $\in \mathbb{R}$

Def: Given  $\psi: F/H \rightarrow U(n)$ , we define

$$P_H^\psi(f) := \rho(N_f, \pi_1(N_f) \rightarrow F/H \xrightarrow{\psi} U(n)) \in \mathbb{R}$$

the Atiyah-Patodi-Singer (APS)  $\rho$ -invrt.

Thm(APS): If  $\psi: \pi_1(N_f) \rightarrow U(n)$  extends to

$\phi: \pi_1(W) \rightarrow U(n)$  then

$$P_H^\psi(f) = \text{sign}^\psi(W) - n \text{sign}(W) \in \mathbb{Z}$$

twisted signature (  $\begin{matrix} \text{signature of finite} \\ \text{Hermitian matrix over } \mathbb{C} \end{matrix}$  )

$$\underline{\text{Ex: }} H = [F, F] \quad \text{and} \quad \pi_1(N_f) \xrightarrow{\psi_H} F/[F, F] = \mathbb{Z}^{2g} \xrightarrow{\phi} U(1)$$

$x_i \longmapsto w_i$

where  $w_i \in \mathbb{C}$  with  $|w_i| = 1$ .

When  $w_i = w$ , define  $\rho_w(f) = \rho(N_f, \pi_1(N_f) \longrightarrow U(1))$

We will show that  $\rho_w(f)$  is often non-trivial.

Note:  $\rho_H^{\psi} : J(H) \rightarrow \mathbb{R}$  is a function. In some cases, it is a homomorphism. However, in general, it is just a "quasi-homomorphism".

Def: A function  $\varphi: G \rightarrow \mathbb{R}$  is called a quasi-homomorphism if  $\exists$  constant  $D(\epsilon)$  such that  $\forall x, y \in G$ ,

$$|\varphi(xy) - \varphi(x) - \varphi(y)| \leq D(\epsilon).$$

$QH(G) := \{ \text{quasi-homomorphisms on } G \} / \text{bounded functions}$

Note: There is an exact sequence

$$0 \rightarrow H^1(G; \mathbb{R}) \rightarrow QH(G) \xrightarrow{\delta} H_b^2(G; \mathbb{R}) \rightarrow H^2(G; \mathbb{R})$$


Proposition (Cochran-H-Horn): The higher p-invs  
 $\rho_H^\psi$  are quasi-homomorphisms.

Idea of Proof ( $\rho_H^*$  is a quasi-homomorphism):

For simplicity, assume  $\Sigma$  is closed.

Then for  $f, h \in J(H)$ , can construct a

4-mfld  $W_{fh}$  by : glue  $M_f \times I$  to  $M_h \times I$   
along  $(\Sigma_g \times (0, 1/2)) \times \{1\} \subset M_f \times \{1\} \cup M_h \times \{1\}$ .



$$\begin{array}{ccc} \Psi_1 : \pi_1 M_f & \rightarrow & F/H \\ \Psi_2 : \pi_1 M_{fg} & \rightarrow & \\ \Psi_3 : \pi_1 M_g & \rightarrow & \end{array}$$

Can show  $\amalg \Psi_i$  extends  $\pi_1(W_{gh}) \rightarrow F/H$ .


One can show that

$$\Rightarrow \left| \rho_H^*(fh) - \rho_H^*(f) - \rho_H^*(h) \right| = \left| \sigma^{(2)}(W, F/H) - \sigma_0(W) \right| \\ \leq 2 \text{ genus } (\Sigma) = D(g)$$

[ $D(g)$  is independent of  $f$  and  $h$ .]

For  $k \geq 1$ , consider


$$P_{w_k}(f) := P(N_f, \pi_1(N_f) \longrightarrow U(1))$$
$$x_i \longmapsto w_k = e^{2\pi i / 4^k}$$



Theorem (Cochran-H-Horn)  $\{P_{w_k}\}_{k=1}^\infty$  is

a linearly independent subset of  $QH(K_g)$  for  $g \geq 2$ .

To prove this, we produce a formula for  
 $P_{W_K}((D_{d_m} \circ D_{B_m})^n)$  for  $k \geq m$  and  $n \geq 0$ .



Ex: For each  $k \geq 2$ , consider  $H = [F_k, F_k]$ .

Since  $[F_k, F_k] \subseteq F_{k+1} = [F, F_k]$ ,

$J([F_k, F_k]) \subseteq J_{k+1}$  ( $k+1^{\text{st}}$  Johnson subgroup).

Def: For  $f \in J(F_k)$ , define

$$\tilde{\rho}_k(f) = \rho^{(2)}(N_f, \pi_1 N_f \rightarrow F/F_k) \in \mathbb{R}.$$

Thm (Cochran - H-Horn): For  $k \geq 2$ ,

$$\tilde{\rho}_k|_{J([F_k, F_k])} : H_1(J([F_k, F_k])) \rightarrow \mathbb{R}$$

is a homomorphism.

Question: What is the image of  $\tilde{\rho}_K$ ? If infinitely generated, then  $H_1(J([F_K, F_K]); \mathbb{Z})$  is infinitely generated.

## Bounded Cohomology

Let  $\rho_H^\psi : J(H) \rightarrow \mathbb{R}$  be one of the  $\rho$ -invs as previously discussed (associated to a finite or infinite unitary representation).

- $\rho_H^\psi : J(H) \rightarrow \mathbb{R}$  is a 1-cochain in group cohomology. Its coboundary

$$\Omega_H^\psi := \delta \rho_H^\psi : J(H) \times J(H) \rightarrow \mathbb{R} \quad \text{is}$$

the (bounded) 2-cocycle

$$\Omega_H^\psi(f, g) := \delta \rho_H^\psi(f, g) = \rho_H^\psi(fg) - \rho_H^\psi(f) - \rho_H^\psi(g).$$

Hence we have the exact sequence :

$$0 \rightarrow H^1(J(H); \mathbb{R}) \rightarrow QH(J(H)) \rightarrow H_b^2(J(H); \mathbb{R})$$
$$\qquad\qquad\qquad \rightarrow H^2(J(H); \mathbb{R})$$

$\overset{\rho_H^*}{\longleftarrow}$

Recall:  $H^1(G; \mathbb{R})$  = homomorphisms  $f: G \rightarrow \mathbb{R}$

$H_b^2(G; \mathbb{R})$  is generated by bounded functions  $f: G \times G \rightarrow \mathbb{R}$  that are cycles

Ex: When  $H = F_3$ ,  $J(H) = K_g$  :

$$0 \rightarrow H^1(K_g; \mathbb{R}) \rightarrow QH(K_g) \xrightarrow{\delta} H_b^2(K_g; \mathbb{R}) \rightarrow \\ R^\infty \stackrel{\psi}{\cong} \{p_{w_K}\} \quad H^2(K_g; \mathbb{R})$$

What is  $\text{im}(\delta|_{\{p_{w_K}\}})$ ,  $\text{ker}(\delta|_{\{p_{w_K}\}})$ ?

Note: If  $\text{ker}(\delta) = R^\infty \Rightarrow \exists \mathbb{Z}^\infty \subset H_1(K_g; \mathbb{Z})$


which answers Morita's question!

## Other Computations of $P^4$

Let  $D \hookrightarrow \Sigma$  be an embedding of

$$D = D^2 - (n\text{-disks})$$

e.g.



$$D \times \{0\} \subset$$


$$\Sigma = 2(D \times I)$$

$P_n$  = pure braid group (framed) is the mapping class group of  $D$ .

This embedding gives an embedding


$$\Theta: P_n \hookrightarrow \text{Mod}(\Sigma).$$

Proposition (Lohran-H-Horn): Let  $\beta \in P_k$ . The higher order  $p$ -invariants of  $\Theta(\beta)$  can be calculated in terms of the higher-order  $p$ -invs of the zero framed surgery on  $\hat{\beta}$ .




## Idea of proof:

$$M_f = \left[ (\Sigma \times S^1) - \left( \text{cylinder diagram} \right) \right] \cup$$



This gives a cobordism  $W$  between



$\Sigma \times S^1$ ,  $M_f$ , and  
 $M_{\hat{\beta}} = 0$ -surgery on  $\hat{\beta}$ .

Signature defect of  $E$  gives the sum  
 of  $\rho$ -invariants on the boundary of  $E$ . B